

Zeros, moments and determinants

Second Symposium on
Analytic Number Theory,
July 9th, 2019

$$\det \begin{pmatrix} \binom{1}{K-1} & \binom{3}{K-1} & \binom{5}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \binom{2}{K-1} & \binom{4}{K-1} & \binom{6}{K-1} & \cdots & \binom{2K}{K-1} \\ \binom{3}{K-1} & \binom{5}{K-1} & \binom{7}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \binom{K}{K-1} & \binom{K+2}{K-1} & \binom{K+4}{K-1} & \cdots & \binom{3K-2}{K-1} \end{pmatrix} = (-1)^{K+1} 2^{\binom{K}{2}}$$

Emilia Alvarez (thesis)



$$\begin{aligned}
& \det_{(K-1) \times (K-1)} \left(\begin{array}{ccccc} \frac{1}{\Gamma(2K-3)} & \frac{1}{\Gamma(2K-5)} & \frac{1}{\Gamma(2K-7)} & \cdots & \frac{1}{\Gamma(1)} \\ \frac{1}{\Gamma(2K-4)} & \frac{1}{\Gamma(2K-6)} & \frac{1}{\Gamma(2K-8)} & \cdots & \frac{1}{\Gamma(0)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{1}{\Gamma(K)} & \frac{1}{\Gamma(K-2)} & \frac{1}{\Gamma(K-4)} & \cdots & \frac{1}{\Gamma(4-K)} \\ \frac{1}{\Gamma(K-1)} & \frac{1}{\Gamma(K-3)} & \frac{1}{\Gamma(K-5)} & \cdots & \frac{1}{\Gamma(3-K)} \end{array} \right) \\
&= \frac{(K-1)(K-2)}{2} \det_{(K-1) \times (K-1)} \left(\begin{array}{ccccc} \frac{1}{\Gamma(2K-3)} & \frac{1}{\Gamma(2K-5)} & \frac{1}{\Gamma(2K-7)} & \cdots & \frac{1}{\Gamma(1)} \\ \frac{1}{\Gamma(2K-4)} & \frac{1}{\Gamma(2K-6)} & \frac{1}{\Gamma(2K-8)} & \cdots & \frac{1}{\Gamma(0)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{1}{\Gamma(K)} & \frac{1}{\Gamma(K-2)} & \frac{1}{\Gamma(K-4)} & \cdots & \frac{1}{\Gamma(4-K)} \\ \frac{1}{\Gamma(K-2)} & \frac{1}{\Gamma(K-4)} & \frac{1}{\Gamma(K-6)} & \cdots & \frac{1}{\Gamma(2-K)} \end{array} \right).
\end{aligned} \tag{1}$$

Ian Cooper (thesis)




$$\begin{vmatrix}
\frac{1}{\Gamma(2k)} & \frac{1}{\Gamma(2k-1)} & \cdots & \frac{1}{\Gamma(k+1)} & \frac{1}{\Gamma(2k)} & \frac{-1}{\Gamma(2k-1)} & \cdots & \frac{(-1)^{k-1}}{\Gamma(k+1)} \\
\frac{1}{\Gamma(2k-1)} & \frac{1}{\Gamma(2k-2)} & \cdots & \frac{1}{\Gamma(k)} & \frac{-1}{\Gamma(2k-1)} & \frac{1}{\Gamma(2k-2)} & \cdots & \frac{(-1)^k}{\Gamma(k)} \\
\vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\
\frac{1}{\Gamma(1)} & \frac{1}{\Gamma(0)} & \cdots & \frac{1}{\Gamma(2-k)} & \frac{-1}{\Gamma(1)} & \frac{1}{\Gamma(0)} & \cdots & \frac{(-1)^{3k-2}}{\Gamma(2-k)}
\end{vmatrix}$$

$$= (-1)^k 2^{k^2} \prod_{\ell=1}^{k-1} \frac{\ell!}{(k+\ell)!}$$

Conrey, Farmer, Keating, Rubinstein, Snaith. Proc. London.
Math. Soc. (3) 91 (2005)




The Riemann Zeta Function



$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad \text{Res} > 1$$
$$= \prod_p (1 - 1/p^s)^{-1}$$

Speiser: RH is equivalent to all complex zeros of $\zeta'(s)$ having real part at least $1/2$



Levinson, Montgomery (1974): $\zeta(s)$ and $\zeta'(s)$ have essentially the same number of zeros in $\Re(s) < 1/2$.

Levinson (1974): More than one third of the zeros of the Riemann zeta function are on the critical line


Conrey (1989): More than two fifths of the zeros of the Riemann zeta function are on the critical line

Bui, Conrey, Young (2011): More than 41% of the zeros of the Riemann zeta function are on the critical line



Horizontal distribution of zeros of the derivative of the Riemann zeta function

Littlewood's lemma (1924): For an analytic function $F(s)$,
(Eg. $\zeta'(s)$)



$$\begin{aligned}
 & \int_T^{T+U} \log |F(b + it)| dt \\
 & - \int_T^{T+U} \log |F(a + it)| dt \\
 & + \int_b^a \arg F(\sigma + i(T + U)) d\sigma \\
 & - \int_b^a \arg F(\sigma + iT) d\sigma \\
 & = 2\pi \sum \text{dist} > 2\pi(a - 1/2) N_L
 \end{aligned}$$

N_L : number of zeros of $\zeta'(s)$ to the left of the $1/2$ -line

$$\int_0^T \log |\zeta'(a + it)| dt$$

$$= \frac{d}{dv} \int_0^T |\zeta'(a + it)|^v dt \Big|_{v=0}$$




Random Unitary Matrices

A is an $N \times N$ unitary matrix:

$$AA^\dagger = A^\dagger A = I$$

eigenvalues of A : $e^{i\theta_n}$


A is chosen randomly with respect to Haar measure on $U(N)$



density of eigenphases: $\frac{N}{2\pi}$

Density of zeros:


$$d(t) \sim \frac{1}{2\pi} \log \frac{t}{2\pi}$$



$$w_n = t_n \frac{1}{2\pi} \log \frac{t_n}{2\pi}, \quad t_n = n^{\text{th}} \text{ Riemann zero}$$


scale the Riemann zeros so that their average spacing is 1

Nearest Neighbor Spacing



Probability density for distances between consecutive eigenvalues/zeros

Using the first 100000 Riemann zeros –
Picture by Andrew Odlyzko




10^5 zeros
around the
 10^{12} th zero



Picture by
A. Odlyzko

billion zeros
around the
 10^{16} th zero



Characteristic polynomial:

$$\begin{aligned}\Lambda_A(s) &= \prod_{n=1}^N (1 - se^{-i\theta_n}) \\ &= \det(I - A^\dagger s)\end{aligned}$$

Equate densities of zeros:


$$\frac{1}{2\pi} \log \frac{\text{RZF}}{T} = \frac{\text{RMT}}{N}$$



Moments of the Riemann Zeta Function

$$\frac{1}{T} \int_0^T |\zeta(1/2 + it)|^2 dt \sim \log T$$

(Hardy and Littlewood, 1918)



$$\frac{1}{T} \int_0^T |\zeta(1/2 + it)|^4 dt \sim \frac{1}{2\pi^2} \log^4 T$$

(Ingham, 1926)

$$\begin{aligned} \zeta(s) &= \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad \text{Res} > 1 \\ &= \prod_p (1 - 1/p^s)^{-1} \end{aligned}$$

Moments of the Riemann Zeta Function: Conjecture

$$\frac{1}{T} \int_0^T |\zeta(1/2 + it)|^{2\lambda} dt \sim a_\lambda f_\lambda \log^{\lambda^2} T$$


Moments of characteristic polynomials: Theorem

$$\int_{U(N)} |\Lambda_A(1)|^{2\lambda} dA_{Haar} \sim f_{\lambda, U(N)} N^{\lambda^2}$$
$$N \sim \log T$$


Conjecture (Keating and Snaith, 2000): $f_\lambda = f_{\lambda, U(N)}$



Zeros of a characteristic polynomial (blue) and its derivative (green)



$N=20$




$N=60$

Pictures by Phan Toan


Horizontal (radial) distribution of zeros of the derivative of zeta (characteristic polynomial)

Mezzadri: J. Phys. A 36 (2003)

Pictures from Dueñez, Farmer, Froelich, Hughes, Mezzadri, Phan: Nonlinearity 23 (2010)



Distribution of distance from the unit circle of zeros of $\Lambda'_A(s)$, $N=100$



Distribution of real part of the zeros of $\zeta'(s)$ for 100000 zeros around $T=1000000$.

Motivated by:

$$\int_0^T |\zeta'(a + it)|^v dt$$

We **would like** to calculate:

$$\int_{U(N)} |\Lambda'_A(s)|^v dA_{Haar} \quad v \text{ non-integer}$$

$v = 1$ Winn: Commun. Math. Phys. 315 (2012)



Motivated by:

$$\int_0^T |\zeta'(a + it)|^v dt$$

We **would like** to calculate:

$$\int_{U(N)} |\Lambda'_A(s)|^v dA_{Haar} \quad v \text{ non-integer}$$

We **can** calculate:

$$\int_{U(N)} |\Lambda'_A(e^{i\theta})|^{2K-2M} |\Lambda_A(e^{i\theta})|^{2M} dA_{Haar}$$

K-M integer

Hughes (2001); Conrey, Rubinstein, Snaith (2006); Dehaye (2008);
Bailey, Bettin, Blower, Conrey, Prokhorov, Rubinstein, Snaith (preprint);
Basor, Bleher, Buckingham, Grava, Its, Its, Keating (preprint)

Motivated by:

$$\int_0^T |\zeta'(a + it)|^v dt$$

We **would like** to calculate:

$$\int_{U(N)} |\Lambda'_A(s)|^v dA_{Haar} \quad v \text{ non-integer}$$

We **can** calculate:

$$\int_{U(N)} \left| \frac{\Lambda'_A}{\Lambda_A}(e^{-\alpha}) \right|^{2K} dA_{Haar}, \quad K \text{ integer}$$

Conrey, Snaith: Commun. Number Th. and Physics. Vol. 2, Num.3 (2008)

Bailey, Bettin, Blower, Conrey, Prokhorov, Rubinstein, Snaith: preprint



Motivated by:

$$\int_0^T |\zeta'(a + it)|^v dt$$

We **would like** to calculate:

$$\int_{U(N)} |\Lambda'_A(s)|^v dA_{Haar} \quad v \text{ non-integer}$$

We **can** calculate:

$$\int_{SO(2N)} (\Lambda''(1))^K dA_{Haar}, \quad K \text{ integer}$$

Altuğ, Bettin, Petrow, Rishikesh, Whitehead (2014)



Motivated by:

$$\int_0^T |\zeta'(a + it)|^v dt$$

We **would like** to calculate:

$$\int_{U(N)} |\Lambda'_A(s)|^v dA_{Haar} \quad v \text{ non-integer}$$

We **can** calculate:

$$\int_{SO(2N)} \left(\frac{\Lambda'_A}{\Lambda_A}(e^{-\alpha}) \right)^K dA_{Haar}, \text{ for integer } K$$

Mason and Snaith (2018)
Emilia Alvarez (thesis)



For large N

$$\int_{SO(2N)} \left(\frac{\Lambda'_A}{\Lambda_A} (e^{-a/N}) \right)^K dA_{Haar}$$
$$\rightarrow \oint \cdots \oint \Delta(u_1^2, u_2^2, \dots, u_K^2) \Delta(u_1, u_2, \dots, u_K) f(u_1) \cdots f(u_K) du_1 \cdots du_K$$

Vandermonde:

$$\Delta(u_1, u_2, \dots, u_K) = \det \begin{vmatrix} 1 & u_1 & u_1^2 & \cdots & u_1^{K-1} \\ 1 & u_2 & u_2^2 & \cdots & u_2^{K-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & u_K & u_K^2 & \cdots & u_K^{K-1} \end{vmatrix}$$



$$\begin{aligned}
& \Delta(u_1^2, u_2^2, \dots, u_K^2) \Delta(u_1, u_2, \dots, u_K) \\
&= \left(\sum_{\sigma \in S_K} \operatorname{sgn}(\sigma) \prod_{i=1}^K u_i^{2\sigma(i)-2} \right) \left(\sum_{\tau \in S_K} \operatorname{sgn}(\tau) \prod_{k=1}^K u_k^{\tau(k)-1} \right) \\
&\rightarrow K! \left(\sum_{\sigma \in S_K} \operatorname{sgn}(\sigma) u_1^{2\sigma(1)-2} u_2^{2\sigma(2)-1} u_3^{2\sigma(3)} \dots u_K^{2\sigma(K)+K-3} \right) \\
&= K! \det(u_i^{2j+i-3})_{i,j=1}^K
\end{aligned}$$



For large N

$$\begin{aligned} & \int_{SO(2N)} \left(\frac{\Lambda'_A}{\Lambda_A} (e^{-a/N}) \right)^K dA_{Haar} \\ & \rightarrow \oint \cdots \oint K! \det(u_i^{2j+i-3})_{i,j=1}^K f(u_1) \cdots f(u_K) du_1 \cdots du_K \\ & \rightarrow K! \det \left(\oint f(u_i) u_i^{2j+i-3} du_i \right)_{i,j=1}^K \end{aligned}$$



$$\begin{aligned}
& \det \begin{pmatrix} \binom{1}{K-1} & \binom{3}{K-1} & \binom{5}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \binom{2}{K-1} & \binom{4}{K-1} & \binom{6}{K-1} & \cdots & \binom{2K}{K-1} \\ \binom{3}{K-1} & \binom{5}{K-1} & \binom{7}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \binom{K}{K-1} & \binom{K+2}{K-1} & \binom{K+4}{K-1} & \cdots & \binom{3K-2}{K-1} \end{pmatrix} \\
&= \binom{n}{r} - \binom{n-1}{r} \\
&= \binom{n-1}{r-1}
\end{aligned}$$

$$\det \begin{pmatrix} \binom{1}{K-1} & \binom{3}{K-1} & \binom{5}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \binom{1}{K-2} & \binom{3}{K-2} & \binom{5}{K-2} & \cdots & \binom{2K-1}{K-2} \\ \binom{2}{K-2} & \binom{4}{K-2} & \binom{6}{K-2} & \cdots & \binom{2K}{K-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \binom{K-1}{K-2} & \binom{K+1}{K-2} & \binom{K+3}{K-2} & \cdots & \binom{3K-1}{K-2} \end{pmatrix}$$

Emilia Alvarez (thesis)



$$\det \begin{pmatrix} \binom{1}{K-1} & \binom{3}{K-1} & \binom{5}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \binom{1}{K-2} & \binom{3}{K-2} & \binom{5}{K-2} & \cdots & \binom{2K-1}{K-2} \\ \binom{1}{K-3} & \binom{3}{K-3} & \binom{5}{K-3} & \cdots & \binom{2K-1}{K-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \binom{1}{1} & \binom{3}{1} & \binom{5}{1} & \cdots & \binom{2K-1}{1} \\ \binom{1}{0} & \binom{3}{0} & \binom{5}{0} & \cdots & \binom{2K-1}{0} \end{pmatrix}$$

$$\binom{2j-1}{1} = 2j - 1$$

$$2 \det \begin{pmatrix} \binom{1}{K-1} & \binom{3}{K-1} & \binom{5}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \binom{1}{K-2} & \binom{3}{K-2} & \binom{5}{K-2} & \cdots & \binom{2K-1}{K-2} \\ \binom{1}{K-3} & \binom{3}{K-3} & \binom{5}{K-3} & \cdots & \binom{2K-1}{K-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \binom{1}{2} & \binom{3}{2} & \binom{5}{2} & \cdots & \binom{2K-1}{2} \\ 1 & 2 & 3 & \cdots & K \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix}$$



$$2 \det \begin{pmatrix} \binom{1}{K-1} & \binom{3}{K-1} & \binom{5}{K-1} & \cdots & \binom{2K-1}{K-1} \\ \binom{1}{K-2} & \binom{3}{K-2} & \binom{5}{K-2} & \cdots & \binom{2K-1}{K-2} \\ \binom{1}{K-3} & \binom{3}{K-3} & \binom{5}{K-3} & \cdots & \binom{2K-1}{K-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \binom{1}{2} & \binom{3}{2} & \binom{5}{2} & \cdots & \binom{2K-1}{2} \\ 1 & 2 & 3 & \cdots & K \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix}$$

$$\binom{2j-1}{2} = 2j^2 - 3j + 1$$

$$\prod_{m=0}^{K-1} \frac{2^m}{m!} \det \begin{pmatrix} 1 & 2^{K-1} & \cdots & K^{K-1} \\ 1 & 2^{K-2} & \cdots & K^{K-2} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 2^2 & \cdots & K^2 \\ 1 & 2 & \cdots & K \\ 1 & 1 & \cdots & 1 \end{pmatrix} = (-1)^{\frac{K(K-1)}{2}} 2^{\binom{K}{2}}$$

